

PACKAGE INSERT**SCHEDULING STATUS:** S0**PROPRIETARY NAME AND DOSAGE FORM:**

Zyzinc+™ Effervescent tablets

COMPOSITION:

Each tablet contains:

Zinc lactate	46,5 mg
providing Zinc (elemental)	10 mg
Vitamin C (ascorbic acid)	1000 mg
Cholecalciferol (Vitamin D3)	(25 µg) 1000 IU
Inactive ingredients: Citric acid, sodium bicarbonate, Sorbitol (sugar alcohol), sodium carbonate, polyethylene glycol, orange flavor, red beet juice powder (colourant), sucralose (artificial sweetener) and riboflavin 5 phosphate sodium (colourant).	
Contains Sorbitol: (sugar alcohol)	
Contains Artificial Sweetener: (Sucralose) 15 mg	
Lactose free, Sugar free, diabetic friendly.	
Suitable for vegetarians.	

PHARMACOLOGICAL CLASSIFICATION:

Category D34.12 Health Supplement – Multiple substance formulation

PHARMACOLOGICAL ACTION:

Vitamin C plays an important role in the body. It is needed to maintain the health of skin, cartilage, teeth, bone and blood vessels. It is also used to protect your body's cells from damage and is known as an antioxidant. Vitamin D contributes to the normal function of the immune system and Zinc is an essential trace element also playing a role in immune function.

INDICATIONS:

Zyzinc+™ is a health supplement for immune support and the maintenance of good health.

CONTRAINDICATIONS:

Individuals with hypersensitivity (allergy) to any of the ingredients should avoid this supplement.

WARNINGS AND SPECIAL PRECAUTIONS:

Do not exceed recommended dosages unless on the advice of a healthcare practitioner. Do not use this product if you are hypersensitive (allergic) to any of the ingredients. If you are on any medication or suffering from any medical condition, it is advisable to seek medical advice before starting any new medicine, supplement or remedy. This unregistered medicine has not been evaluated by the South African Health Products Regulatory Authority for its quality, safety or intended use. Health supplements are intended only to complement health or supplement the diet.

INTERACTIONS:

High doses of Vitamin C may reduce the response to warfarin, possibly by causing diarrhea and reducing warfarin absorption. Patients to consult their Oncologist before using Vitamin C supplements, especially in high doses. Taking Vitamin D along with digoxin might increase the effects of digoxin and lead to an irregular heartbeat. The use of Vitamin D along with diltiazem might decrease the effectiveness of diltiazem. The use of Vitamin D with thiazide diuretics may contribute to hypercalcaemia. The concomitant use of Zinc reduces the absorption of penicillamine. The concomitant use of Zinc also reduces the absorption of quinolone and tetracycline antibiotics.

PREGNANCY AND LACTATION:

Safety for use during pregnancy and breastfeeding has not been established.

DOSAGE AND DIRECTIONS FOR USE:

Adults and children over 9 years: Take 1 (one) tablet daily dissolved in half to a full glass of water.

SIDE EFFECTS:

This product may cause mild gastrointestinal disturbances.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

Treatment should be symptomatic and supportive. In the case of accidental overdose of the product, contact the nearest hospital or poison control centre.

IDENTIFICATION:

Light pink to beige round flat tablet, with a bevelled edge.

PRESENTATION:

Zyzinc+™ effervescent tablets are packed in polypropylene tubes of 20's and packed into a carton.

STORAGE INSTRUCTIONS:

Store in a cool dry place below 25 °C, protect from light.

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER: Not applicable.

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION:

Sunstone Brands (Pty) Ltd
1073 Anvil Street
Robertville
Florida
1710

DATE OF PUBLICATION OF THE PACKAGE INSERT:

February 2021

ZYL-335

VOUBILJET**SKEDULERINGSTATUS:** S0**HANDELSNAAM EN DOSEERVORM:**

Zyzinc+™ Bruistablette

SAMESTELLING:

Elke bruistablet bevat:

Sink laktat	46,5 mg
verskaf Sink (elementele)	10 mg
Vitamiën C (askorbiensuur)	1000 mg
Cholekalsiferol (Vitamiën D)	(25 µg) 1000 IU
Onaktiewe bestanddele: Sitroensuur, natriumbikarbonaat, Sorbitol (suiker alkohol), natriumkarbonaat, poliëteleen glikol, lemoen geursel, Beetsap poeier (kleursidant), sukralose (kunsmatige versoeter) en riboflavien 5 fosfaat natrium (kleursel).	
Laktosevry, suikervry en geskik vir diabetie.	
Geskik vir vegetariërs.	

FARMAKOLOGIESE KLASSIFIKASIE:

Kategorie D34.12. Gesondheidsaanvulling – Meervoudige stof samestelling

FARMAKOLOGIESE AKSIE:

Vitamiën C speel 'n belangrike rol in die liggaam. Dit is nodig om die gesondheid van vel, kraakbeen, tande, been en bloedvate te handhaaf. Dit word ook gebruik om liggaamselle teen skade te beskerm en staan bekend as 'n antioksidant. Vitamiën D dra by tot die normale funksie van die immuunstelsel en Sink is 'n essensiële sporelement wat ook 'n rol speel in immuunfunksie.

INDIKASIES:

Zyzinc+™ is 'n gesondheidsaanvulling vir immuun ondersteuning en die handhawing van goeie gesondheid.

KONTRAINDIKASIES:

Individue met hipersensitiwiteit (allergie) vir enige van die bestanddele moet hierdie aanvulling vermy.

WAARSKUWINGS EN SPESIALE VOORSORGMAATREËLS:

Moenie die voorgestelde dosis oorskry nie tensy aanbeveel deur 'n gesondheidsorgpraktisyn. Moenie hierdie produk gebruik as jy hipersensitief (allergies) is vir enige van die bestanddele nie. Indien jy enige ander medikasie gebruik of ly aan 'n mediese kondisie, is dit raadsaam om eers mediese advies te bekom voor enige nuwe medikasie, aanvullings of gesondheidsmiddels gebruik word. Hierdie ongeregistreerde medisyne is nie deur die Suid Afrikaanse Gesondheidsprodukte Regulatoriese Owerheid geëvalueer vir kwaliteit, veiligheid of die beoogde gebruik nie. Gesondheidsaanvullings is slegs bedoel om gesondheid te komplementeer of die dieet aan te vul.

INTERAKSIES:

Hoë dosisse Vitamiën C kan die reaksie op warfarin verminder, moontlik deur diarree te veroorsaak en die opname van warfarin te verminder. pasiënte moet hul Onkoloog raadpleeg voordat hulle Vitamiën C aanvullings gebruik, veral in hoë dosisse. As Vitamiën D saam met digoxin geneem word, kan dit die effek van digoxin verhoog en dit kan lei tot 'n onreëlmatige hartklop. Die gebruik van Vitamiën D saam met diltiazem kan die effektiwiteit van diltiazem verminder. Die gebruik van Vitamiën D saam met tiazied-diuretika kan bydra tot hiperkalsemie. Die gepaardgaande gebruik van Sink verminder die opname van penisillamien. Die gepaardgaande gebruik van Sink verminder ook die opname van kinolon- en tetrasiklienantibiotika.

SWANGERSKAP EN LAKTASIE:

Veiligheid vir gebruik gedurende swangerskap en laktasie is nie vasgestel nie.

DOSIS EN GEBRUIKSAANWYSINGS:

Volwassenes en kinders ouer as 9 jaar: Neem 1 (een) tablet daagliks opgelos in 'n halwe of vol glas water.

NEWE-EFFEKTE:

Hierdie produk mag lei tot ligte versteurings van die spysverteringskanaal.

BEKENDE SIMPTOME VAN OORDOSERING EN**BESONDERHEDE VIR DIE BEHANDELING DAARVAN:**

Behandeling moet simptomaties en ondersteunend wees. In geval van onopsettlike oordosering van die produk, kontak die naaste hospitaal of gifbeheersentrum.

IDENTIFIKASIE:

Ligte pienk tot rooimkleur ronde plat tablet, met 'n afgeskuinste rand.

AANBIEDING:

Zyzinc+™ Bruistablette is verpak in polipropileen buise van 20's en verpak in 'n karton.

BERGINGSANWYSINGS:

Bewaar in 'n koel, droë plek benede 25 °C, beskerm teen lig.

HOU BUITE BEREIK VAN KINDERS.

REGISTRASIE-NOMMER: Nie van toepassing.

NAAM EN BESIGHEIDSDRES VAN DIE APPLIKANT VAN**DE DIE REGISTRASIEERTIFIKAAT:**

Sunstone Brands (Pty) Ltd
1073 Anvil Street
Robertville
Florida
1710

DATUM VAN PUBLIKASIE VAN DIE VOUBILJET:

Februarie 2021

ZYL-335

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S0

PROPRIETARY NAME AND DOSAGE FORM:

Zyzinc+™ Effervescent tablets

Read all of this leaflet carefully because it contains important information for you. Zyzinc+™ is available without a doctor's prescription and is a complementary medicine. Nevertheless you still need to use zyzinc+™ carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share zyzinc+™ with any other person.
- Ask your pharmacist or healthcare practitioner if you need more information or advice.

1. WHAT ZYZINC+™ CONTAINS:

Each tablet contains the following active materials:

Zinc lactate 46,5 mg (providing Zinc 10 mg), Ascorbic Acid 1000 mg (providing Vitamin C 1000 mg) and Vitamin D 1000 IU.

Inactive ingredients: Citric acid, sodium bicarbonate, Sorbitol (sugar alcohol), sodium carbonate, polyethylene glycol, orange flavor, red beet juice powder (colourant), sucralose (artificial sweetener) and riboflavin 5 phosphate sodium (colourant).

Contains Sorbitol: (sugar alcohol)

Contains Artificial Sweetener: (Sucralose) 15 mg

Lactose free, Sugar free, diabetic friendly.

Suitable for vegetarians.

2. WHAT ZYZINC+™ IS USED FOR:

Zyzinc+™ is a health supplement for immune support and the maintenance of good health.

3. BEFORE YOU TAKE ZYZINC+™:

Consult a healthcare practitioner prior to use if you are taking warfarin. Consult with your Oncologist before taking this supplement. Also consult with your healthcare practitioner prior to use if you are on digoxin, diltiazem, thiazide diuretics, quinolone or tetracycline antibiotics.

Do not take zyzinc+™ if you are hypersensitive (allergic) to any of the ingredients. This unregistered medicine has not been evaluated by the South African Health Products Regulatory Authority for its quality, safety or intended use. Health supplements are intended only to complement health or supplement the diet.

PREGNANCY AND BREASTFEEDING:

Safety for use during pregnancy and breastfeeding has not been established.

4. HOW TO TAKE ZYZINC+™:

Adults and children over 9 years: Take 1 (one) tablet daily dissolved in half to a full glass of water.

IF YOU TAKE MORE ZYZINC+™ THAN YOU SHOULD:

In the event of overdose, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre.

5. POSSIBLE SIDE EFFECTS:

Side effects may include mild gastrointestinal disturbances. Not all side effects reported for zyzinc+™ are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking this medicine, consult your doctor, pharmacist or other healthcare professional for advice.

6. STORING AND DISPOSING OF ZYZINC+™:

- Keep all medicines out of reach and sight of children
- Store at or below 25 °C
- Protect from light and moisture
- Do not store in the bathroom
- Store in its original container
- Keep the container tightly closed
- Do not use after the expiry date stated on the container
- Do not use zyzinc+™ if you notice visible signs of deterioration
- Return all unused medicine to your pharmacist
- Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

7. PRESENTATION OF ZYZINC+™:

Zyzinc+™ effervescent tablets are packed in polypropylene tubes of 20's and packed into a carton.

8. IDENTIFICATION OF ZYZINC+™:

Light pink to beige round flat tablet, with a bevelled edge.

9. REGISTRATION NUMBER:

Registry number: Not applicable

10. NAME AND ADDRESS OF REGISTRATION HOLDER:

Sunstone Brands (Pty) Ltd
1073 Anvil Street
Robertville
Florida
1710

11. DATE OF PUBLICATION:

February 2021
ZYL3-335

PASIEËNTINLIGTINGSBLAADJIE

SKEDULERINGSSTATUS: S0

HANDELSNAAM EN DOSEERVORM:

Zyzinc+™ Bruistablette

Lees die hele inligtingsblaadjie noukeurig deur, dit bevat belangrike inligting vir jou. Zyzinc+™ is beskikbaar sonder 'n doktersvoorskrif en is aanvallende (komplimentêre) medisyne. Jy moet egter nogsteeds zyzinc+™ versigtig gebruik om die beste resultate te verkry.

- Hou hierdie voulijet, dit mag dalk nodig wees dat jy dit weer lees.
- Moenie zyzinc+™ deel met enige ander persoon nie.
- Raadpleeg jou apteker of gesondheidsorgpraktisyn indien jy meer inligting of advies benodig.

1. WAT ZYZINC+™ BEVAT:

Elke tablet bevat die volgende aktiewe bestanddele:

Sink laktaat 46,5 mg (verskaf Sink 10 mg), Ascorbiensuur 1000 mg (verskaf Vitamien C 1000 mg) en Vitamien D 1000 IU.

Onaktiewe bestanddele: Sitroensuur, natriumbikarbonaat, Sorbitol (suiker alkohol), natriumkarbonaat, poliëteleen glikol, lemoen geursel, Beetsap poeier (kleursel), sukralose (kunsmatige versoeter) en riboflavin 5 fosfaat natrium (kleursel).

Laktosevry, suikervry en geskik vir diabetie.

Geskik vir vegetariërs.

2. WAARVOOR ZYZINC+™ GEBRUIK WORD:

Zyzinc+™ is 'n gesondheidsaanvulling vir die ondersteuning van gesonde vel, hare en naels.

3. VOORDAT ZYZINC+™ GEBRUIK WORD:

Raadpleeg jou gesondheidsorgpraktisyn voor gebruik indien jy warfarin neem. Raadpleeg jou Onkoloog voor die gebruik van hierdie aanvulling. Raadpleeg ook jou Gesondheidsorgpraktisyn voor gebruik indien jy digoxin, diltiazem, tiazied diuretika, kinolon of tetrakislienantibiotika neem.

Moenie zyzinc+™ gebruik as jy hipersensitief (allergies) is vir enige van die bestanddele nie. Hierdie ongeregistreerde medisyne is nie deur die Suid Afrikaanse Gesondheidsprodukte Regulatoriese Owerheid geëvalueer vir kwaliteit, veiligheid of die beoogde gebruik nie. Gesondheidsaanvullings is slegs bedoel om gesondheid te komplementeer of die dieet aan te vul.

SWANGERSKAP EN BORSVOEDING:

Veiligheid vir gebruik gedurende swangerskap en laktasie is nie vasgestel nie.

4. HOE OM ZYZINC+™ TE GEBRUIK:

Volwassenes en kinders ouer as 9 jaar: Neem 1 (een) tablet daagliks opgelos in 'n halwe of vol glas water.

INDIEN JY MEER ZYZINC+™ GEBRUIK AS WAT JY BEHOORT:

Indien oordosering vermoed word, raadpleeg onmiddellik jou dokter of apteker. Indien geneeën beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum.

5. MOONTLIKE NEWE-EFFEKTE:

Nuwe-effekte kan simptome van 'n omgekrapte spysverteringskanaal insluit. Nie alle nuwe-effekte gerapporteer vir zyzinc+™ is ingesluit in hierdie pamflet nie. Indien jou algemene gesondheid versleg of jy enige ongewenste effekte ervaar terwyl hierdie medisyne geneem word, raadpleeg jou dokter, apteker of ander gesondheidsorgpraktisyn.

6. BERGING EN WEGDOEN VAN ZYZINC+™:

Bewaar alle medikasie buite bereik en sig van kinders

- Bewaar teen of benede 25 °C
- Beskerm teen lig en vog
- Moenie die produk in die badkamer bewaar nie
- Bewaar in die oorspronklike houer
- Hou die houer dig toe
- Moenie gebruik na die vervaldatum aangedui op die houer nie
- Moenie zyzinc+™ gebruik indien jy enige sigbare tekens van produkafbraak opmerk nie.
- Besorg enige ongebruikte medisyne terug aan jou apteker.
- Moenie van ongebruikte medisyne ontslae raak in dreine of rioolstelsels nie (bv. toilette).

7. AANBIEDING VAN ZYZINC+™:

Zyzinc+™ Bruistablette is verpak in polipropileen buise van 20's en verpak in 'n karton.

8. IDENTIFIKASIE VAN ZYZINC+™:

Ligte pienk tot rooimkleur ronde plat tablet, met 'n afgeskuinste rand.

9. REGISTRASIONOMMER:

Registrasienommer: Nie van toepassing.

10. NAAM EN BESIGHEIDSADRES VAN DIE APPLIKANT VAN DIE REGISTRASIESERTIFIKAAT:

Sunstone Brands (Pty) Ltd
1073 Anvil Straat
Robertville
Florida
1710

11. DATUM VAN PUBLIKASIE:

Februarie 2021
ZYL3-335